

The causal ladder and the strength of K -causality. II

E. Minguzzi *

Abstract

Hawking's stable causality implies Sorkin and Woolgar's K -causality. The work investigates the possible equivalence between the two causality requirements, an issue which was first considered by H. Seifert and then raised again by R. Low after the introduction of K -causality. First, a new proof is given that a spacetime is stably causal iff the Seifert causal relation is a partial order. It is then shown that given a K -causal spacetime and chosen an event, the light cones can be widened in a neighborhood of the event without spoiling K -causality. The idea is that this widening of the light cones can be continued leading to a global one. Unfortunately, due to some difficulties in the inductive process the author was not able to complete the program for a proof as originally conceived by H. Seifert. Nevertheless, it is proved that if K -causality coincides with stable causality then in any K -causal spacetime the K^+ future coincides with the Seifert future. Explicit examples are provided which show that the K^+ future may differ from the Seifert future in causal spacetimes.

1 Introduction

The property of K -causality was introduced about ten years ago by R.D. Sorkin and E. Woolgar [11]. Given a spacetime (M, g) they defined the relation K^+ as the smallest closed subset $K^+ \subset M \times M$, which contains I^+ , $I^+ \subset K^+$, and shares the transitivity property: $(x, y) \in K^+$ and $(y, z) \in K^+ \Rightarrow (x, z) \in K^+$ (the set of causal relations satisfying these properties is non-empty, consider for instance the trivial subset $M \times M$). This definition raised from the fact that J^+ while transitive is not necessarily closed whereas $\bar{J}^+$ while closed is not necessarily transitive. They also defined the spacetime to be K -causal if the relation K^+ is a partial order i.e. not only transitive and reflexive but also antisymmetric, that is, such that, $(x, z) \in K^+$ and $(z, x) \in K^+ \Rightarrow x = z$.

R. Low pointed out [11, footnote p. 1990] that H. Seifert's causal relation $J_S^+ \supset J^+$ is closed and transitive, hence $K^+ \subset J_S^+$. It is then natural to ask whether it is always $K^+ = J_S^+$ and if this is not the case, whether it is at least

*Department of Applied Mathematics, Florence University, Via S. Marta 3, 50139 Florence, Italy. E-mail: ettore.minguzzi@unifi.it

true that J_S^+ is a partial order whenever K^+ is a partial order. Seifert proved [10] the transitivity and closure of J_S^+ and gave an argument showing that J_S^+ is a partial order if and only if the spacetime is stably causal (for a rigorous proof see [4, Prop. 2.3] or theorem 3.12 below). As a consequence, since $K^+ \subset J_S^+$, if the spacetime is stably causal then it is K -causal. Moreover, the equality $K^+ = J_S^+$ would imply that the properties of K -causality and stable causality coincide. On the contrary, the example of a spacetime K -causal but non-stably causal would imply at once that K^+ does not always coincide with J_S^+ and that K -causality can be included in the causal hierarchy of spacetimes [4, 6] just below stable causality.

Seifert himself [10] raised the problem as to whether J_S^+ could be regarded as the smallest closed and transitive causal relation containing I^+ . One of his lemmas [10, Lemma 2] actually answers this question in the affirmative sense provided K^+ is a partial order.

Claim 1.1. (*Seifert's [10, Lemma 2]*)

J_S^+ is the smallest among the partial orders $P^+ \subset M \times M$ not smaller than J^+ with closed $P^+(x)$ and $P^-(x)$ for all x [provided such a smallest partial order exists]¹.

Indeed, we shall see in section 2 that the previous claim can be conveniently rephrased in the following way

Claim 1.2. *If (M, g) is K -causal then $K^+ = J_S^+$.*

A consequence is that K -causality is equivalent to stable causality. Unfortunately, Seifert's arguments were not rigorous as they did not take into account the many subtleties of the K^+ relation. This lemma was almost never cited in subsequent literature and some researchers who tried to reproduce it began to raise some doubts on its validity. It suffices to say that using it some proofs later given by Hawking and Sachs [4] could have been considerably simplified as I will show in the last section. In that fundamental work Hawking and Sachs preferred to take a path independent of Seifert's 1971 work and indeed, although they cited Seifert, they gave a completely new proof of the equivalence between stable causality and the antisymmetry of J_S^+ and avoided any mention to the claim above. Over the time the question raised by Seifert's work was overlooked and only with the introduction of the K^+ relation it was rediscovered from a different perspective. This work is devoted to the study of this open issue.

The work is organized as follows.

In section 2 some general results for binary relations on $M \times M$ are given. The equivalence between claims 1.1 and 1.2 is proved here. In this section as well as in the rest of the work the reader is assumed to be familiar with the conventions and notations introduced in [5, Sect. 1 and 2]. Let me just remind that the spacetime signature is $(-, +, \dots, +)$, that the subset inclusion is reflexive, $X \subset X$, and that the boundary of a set A is denoted $\bar{A}$.

¹Text in square brackets by the author.

Section 3 deals with Seifert's closed relation. I generalize, simplify and fill gaps of some proofs, particularly that on the equivalence between stable causality and the antisymmetry of J_S^+ , the extent of the improvements being there explained.

Section 4 deals with some results on the violating sets for J_S^+ .

In section 5 some examples of spacetimes in which $K^+ \neq J_S^+$ are given. The strategy outlined by Seifert for a proof of the equivalence between K -causality and stable causality is, more or less, followed here and made rigorous. Seifert suggested to prove that (i) K -causality implies strong K -causality ([11, Lemma 16] and lemma 5.5 below), and that (ii) strong K -causality implies that, chosen an event, the cones can be widened in a neighborhood of the chosen point while preserving K -causality (theorem 5.19) (iii) the process of widening the light cones can be continued so as to obtain a global widening of the light cones and hence the proof that a K -causal spacetime is stably K -causal and hence stably causal (actually stable causality and stable K -causality coincide, see corollary 2.3).

The same strategy is followed here although in several points the proofs differ significantly from what originally suspected by Seifert. Several technical lemmas are required because what is intuitive for J^+ is usually hard to prove for K^+ , the reason lying on the fact that K^+ is defined through its closure and transitivity properties and not, at least not straightforwardly, by using the set of causal curves. Indeed, Sorkin and Woolgar [11] introduced the method of transfinite induction in order to obtain some basic results. Here it is shown that this method can be avoided, and that it can be replaced by soft topological arguments which take advantage of the minimality of K^+ .

Unfortunately, I will not be able to prove step (iii). The process of widening the light cones can indeed be indefinitely continued but some technicalities do not allow to conclude that a global widening of the light cones can be obtained.

Nevertheless, in section 6 it is proved that if it is true that K -causality coincides with stable causality then it is also true that in a K -causal spacetime $K^+ = J_S^+$ (theorem 6.2). Under the same assumption a very simple proof that causal continuity implies stable causality is given. In this respect the proof of the equivalence between K -causality and stable causality is recognized as an important problem in causality theory as many old proof would be greatly simplified by the knowledge of this result, and new ones would follow.

Despite the fact that the main problem has not been solved, the sections 5 and 6 contain many new results and properties of the K^+ relation which may prove to be useful in future applications.

2 Preliminaries

The reader is assumed to be familiar with [5, Sect. 2] which contains some definitions and results for a generic binary relation $R^+ \subset M \times M$. The definitions of closure, partial closure, transitivity, reflexivity, antisymmetry, R -causality, strong R -causality, stable R -causality will not be repeated here. Nor will be

repeated the definitions of the sets $R^\pm(x)$ or R^- given R^+ or of the diagonal Δ in $M \times M$.

Here I give some useful definitions for dealing with violations of the antisymmetry condition. The *R-causality violating set on $M \times M$* is

$$VR = R^+ \cap R^- \cap \Delta^C, \quad (1)$$

which is the set of pairs at which the antisymmetric condition for R^+ (and hence R^-) fails. Note that since Δ is closed VR is open whenever R^+ is open, for instance, VI is open.

Analogously, it is useful to define the *R-causality violating set on M* as $vR = \pi(VR)$, where $\pi : M \times M \rightarrow M$ is the projection on the first factor (or, equivalently, on the second factor). It is made of all the points x at which *R-causality* at x does not hold.

In general given the spacetime (M, g) the causal relation R^+ will be related in some way to (M, g) , for instance I^+ is the set of pairs (x, z) such that there is a timelike curve connecting x to z . In order to stress this spacetime dependence I shall sometimes write $R_{(M, g)}^+$. If we are working in the same spacetime manifold M but with different metrics I may write R_g^+ in place of $R_{(M, g)}^+$ in order to stress the metric (or most often the conformal structure) dependence. Analogously I may write sentences like “ $(\bar{g})$ -causal curve γ ” to stress that γ is causal with respect to the metric $\bar{g}$, or “ $(\bar{g})$ -convex set V ” to stress that V is a convex set with respect to the metric $\bar{g}$. If instead we are dealing with the same metric but with different open sets $A \subset M$, I may write R_A^+ in place of $R_{(A, g)}^+$. However, the reader should be careful because J_S^+ denotes the Seifert future, not the causal relation for the spacetime (S, g) . In order to avoid confusion I will not use the letter S for any open set.

Let me recall [5, Th. 2.2] that a transitive relation R^+ which contains I^+ is closed iff it has closed $R^\pm(x)$ for all x , and that, as a consequence [5, Cor. 2.5], K^+ is the smallest transitive relation containing I^+ such that for every $x \in M$, $K^+(x)$ and $K^-(x)$ are closed. Theorem 2.2 of [5] implies

Corollary 2.1. *Claims 1.1 and 1.2 are equivalent.*

Proof. Note that the smallest relation, call it M^+ , which is transitive, contains I^+ , is partially closed (or equivalently closed because of the first two properties) and antisymmetric may not exist because though all these properties are preserved under arbitrary intersections of relations, the set of relations satisfying the properties may be empty. Provided it is not empty M^+ exists and $K^+ \subset M^+$ because M^+ is transitive, closed, and contains I^+ . But then the inclusion implies that K^+ is antisymmetric too thus the spacetime is K -causal, and by definition of M^+ , $M^+ \subset K^+$. Thus if M^+ exists $M^+ = K^+$ and the spacetime is K -causal. Conversely, if the spacetime is K -causal then M^+ exists and hence $M^+ = K^+$. As a consequence, claims 1.1 and 1.2 are equivalent. $\square$

Let $\mathcal{P}$ be a conformal invariant property for a spacetime. Assume moreover that if $\mathcal{P}$ holds for (M, g_1) then it holds for every (M, g_2) , $g_2 < g_1$. Examples

for $\mathcal{P}$ are chronology, causality, distinction, K -causality and stable causality. For such a property the spacetime (M, g) is said to have the *stable- $\mathcal{P}$ property* or to be *stably $\mathcal{P}$* if there is $g' > g$ such that (M, g') has the $\mathcal{P}$ property. It is clear that stable-causality is the usual stable causality. It is also clear that if $\mathcal{P}_1 \Rightarrow \mathcal{P}_2$ then $\text{stable-}\mathcal{P}_1 \Rightarrow \text{stable-}\mathcal{P}_2$. A nice result is that the operation of making a property stable is idempotent, that is

Lemma 2.2. *Let $\mathcal{P}$ be a conformal invariant property such that if $\mathcal{P}$ holds for (M, g_1) then it holds for every (M, g_2) , $g_2 < g_1$. Then the property stable-stable- $\mathcal{P}$ coincides with stable- $\mathcal{P}$.*

Proof. The implication $\Rightarrow$ is obvious, for the other direction if (M, g) is stably $\mathcal{P}$ there is $g' > g$ such that (M, g') has property $\mathcal{P}$. Take g'' , $g < g'' < g'$, then (M, g'') is stably $\mathcal{P}$ and hence (M, g) has property stable-stable- $\mathcal{P}$. $\square$

Corollary 2.3. *Stable causality is equivalent to stable chronology, stable distinction, stable strong causality, stable K -causality and stable stable causality.*

Proof. Stable chronology implies causality, because a closed causal curve for (M, g) is closed and timelike for (M, g') , $g' > g$. Thus stable-chronology = stable-stable-chronology $\Rightarrow$ stable-causality. Now consider the closed chain of implications stable-chronology $\Rightarrow$ stable-causality = stable-stable-causality $\Rightarrow$ stable- K -causality $\Rightarrow$ stable-strong causality $\Rightarrow$ stable-distinction $\Rightarrow$ stable-causality $\Rightarrow$ stable chronology, from which the equivalence of all these properties follows. $\square$

Stable non-total viciousness is distinct from stable causality (see remark 4.5).

3 Seifert's closed relation and stable causality

In this section the relationship between stable causality and Seifert's causal relation J_S^+ is clarified. Concerning the topology of stable causality, for some results not considered here nor cited elsewhere in the paper but still of interest the reader may consult [1, 9] and [3, Sect. 6.4].

Given two metrics g, g' , over M , denote as usual $g' > g$ if every causal vector for g is timelike for g' , and $g' \geq g$ if every causal vector for g is causal for g' . In presence of different metrics, the sets $I_g^+, J_g^+ \subset M \times M$, are the chronological and causal sets of (M, g) .

Definition 3.1. The set $J_S^+ \subset M \times M$, defining Seifert's causal relation on (M, g) is

$$J_S^+ = \{(x, z) : (x, z) \in J_{g'}^+ \text{ for every } g' > g\} = \bigcap_{g' > g} J_{g'}^+.$$

Sometimes, in order to avoid confusion, I shall write $J_{S_g}^+$ in place of J_S^+ to point out the causal structure to which J_S^+ refers to.

Lemma 3.2. *If $\tilde{g} < g$ then $\bar{J}_{\tilde{g}}^+ \subset \Delta \cup I_g^+$.*

Proof. Let $(x, z) \in \bar{J}_{\bar{g}}^+ \setminus \Delta$, let σ_n be a sequence of $(\tilde{g})$ -causal curves of endpoints x_n, z_n in $(M, \tilde{g})$. Using a limit curve theorem it follows the existence of a future directed $(\tilde{g})$ -causal curve σ^x starting from x , and a past directed $(\tilde{g})$ -causal curve σ^z ending at z , and a subsequence σ_j distinguishing both curves. Taken $x' \in \sigma^x$, $z' \in \sigma^z$ it follows $(x, x') \in J_{\bar{g}}^+$, $(z', z) \in J_{\bar{g}}^+$ and $(x', z') \in \bar{J}_{\bar{g}}^+$, or, in terms of the causal relations of g , $(x, x') \in I_g^+$, $(z', z) \in I_g^+$ and $(x', z') \in \bar{J}_g^+$, which implies, because I_g^+ is open, $(x, z) \in I_g^+$. $\square$

Lemma 3.3. *Equivalent definitions of J_{Sg}^+ on the spacetime (M, g) are*

$$J_{Sg}^+ = \Delta \cup \bigcap_{g' > g} I_{g'}^+ = \bigcap_{g' > g} \bar{J}_{g'}^+,$$

moreover, $J_{Sg}^+ = \bigcap_{g' > g} J_{Sg'}^+$.

Proof. For the first equality we have only to show that $\bigcap_{g' > g} J_{g'}^+ \subset \Delta \cup \bigcap_{g' > g} I_{g'}^+$ the other inclusion being obvious. For every $\bar{g} > g$, taken $\tilde{g}$ such that $g < \tilde{g} < \bar{g}$, it is $J_{\tilde{g}}^+ \subset \Delta \cup I_{\tilde{g}}^+$, hence $\bigcap_{g' > g} J_{g'}^+ \subset \Delta \cup I_{\tilde{g}}^+$, since $\bar{g} > g$ is arbitrary the thesis follows.

For the second equality we have only to show that $\bigcap_{g' > g} \bar{J}_{g'}^+ \subset \bigcap_{g' > g} J_{g'}^+$ the other inclusion being obvious. Let $\bar{g} > g$, taken $\tilde{g}$ such that $g < \tilde{g} < \bar{g}$, by lemma 3.2 it is $\bar{J}_{\tilde{g}}^+ \subset J_{\tilde{g}}^+$, thus $\bigcap_{g' > g} \bar{J}_{g'}^+ \subset J_{\tilde{g}}^+$, since $\bar{g} > g$ is arbitrary the thesis follows.

For the last statement note that $J_{Sg}^+ = \bigcap_{g' > g} J_{g'}^+ = \bigcap_{g' > g} (\bigcap_{g'' > g'} J_{g''}^+) = \bigcap_{g' > g} J_{Sg'}^+$. $\square$

Theorem 3.4. *The relation J_S^+ is closed, transitive and contains I^+ , moreover for every $x \in M$, $J_S^+(x)$ and $J_S^-(x)$ are closed and contain respectively $I^+(x)$ and $I^-(x)$.*

Proof. The transitivity is obvious because for every $g' > g$, $J_{g'}^+$ is transitive. The statements on the inclusions of I^+ , $I^+(x)$ and $I^-(x)$ are trivial too. The closure of J_S^+ is immediate from $J_S^+ = \bigcap_{g' > g} \bar{J}_{g'}^+$. The closure of $J_S^-(x)$, and $J_S^+(x)$, follows by taking $(x_n, z_n) \rightarrow (x, z)$, $(x_n, z_n) \in J_S^+$, and by using the closure of J_S^+ in the cases $x_n = x$ or $z_n = z$ for the initial choice of converging sequence. $\square$

A simple consequence is $K^+ \subset J_S^+$ and hence

Corollary 3.5. *If J_S^+ is a partial order then the spacetime is K -causal.*

Lemma 3.6. *If $g < g'$ then $K_g^+ \subset \Delta \cup I_{g'}^+$.*

Proof. It follows from $K_g^+ \subset J_{Sg}^+ = \Delta \cup \bigcap_{g' > g} I_{g'}^+ = \bigcap_{g' > g} (\Delta \cup I_{g'}^+)$. $\square$

Lemma 3.7. *An equivalent definition of $J_{S_g}^+$ on the spacetime (M, g) is*

$$J_{S_g}^+ = \bigcap_{g' > g} K_{g'}^+.$$

Proof. Since $J_{g'}^+ \subset K_{g'}^+$, $J_{S_g}^+ = \bigcap_{g' > g} J_{g'}^+ \subset \bigcap_{g' > g} K_{g'}^+$, but, since $J_{S_{g'}}^+$ is closed and transitive, $K_{g'}^+ \subset J_{S_{g'}}^+$ and $\bigcap_{g' > g} K_{g'}^+ \subset \bigcap_{g' > g} J_{S_{g'}}^+ = J_{S_g}^+$, from which the thesis follows. $\square$

Remark 3.8. From the definition of the sets $R^\pm(x)$ given the binary relation R^+ it follows

$$J_{S_g}^\pm(x) = \{x\} \cup \bigcap_{g' > g} I_{g'}^\pm(x) = \bigcap_{g' > g} J_{g'}^\pm(x) = \bigcap_{g' > g} K_{g'}^\pm(x) = \bigcap_{g' > g} J_{S_{g'}}^\pm(x).$$

Lemma 3.9. *If J_S^+ on (M, g) is a partial order then for every $x \in M$ there is a $(x$ -dependent) metric $g_x > g$ such that (M, g_x) is chronological at x .*

Proof. Assume, by contradiction, that the thesis does not hold, then there is $x \in M$ such that for every $g' > g$, there is a closed (g') -timelike curve $\sigma_{g'}$ passing through x .

Let $\bar{g} > g$, introduce a Riemannian metric in a neighborhood of x and consider $S = \bar{B}(x, \epsilon)$, i.e. the surface of the ball of Riemannian radius $\epsilon > 0$. Choose ϵ sufficiently small so that S is contained in a $(\bar{g})$ -convex neighborhood contained in a $(\bar{g})$ -globally hyperbolic neighborhood V .

For every g' , $g < g' < \bar{g}$, there is a closed (g') -timelike curve $\sigma_{g'}$ passing through x . This curve must escape the hyperbolic neighborhood V otherwise in $(V, \bar{g})$ there would be a closed $(\bar{g})$ -timelike curve. Hence the curve must meet S at some point of $S \cap J_{\bar{g}}^+(x)$. Given g' the event x belongs to the *chronologically violating set* $vI'_{g'}$ which is open [8] and which can be written as the union of disjoint open sets of the form $I_{g'}^+(y) \cap I_{g'}^-(y)$ where y is any point of the component [8, Prop. 4.27]. In particular x belongs to the component $I_{g'}^+(x) \cap I_{g'}^-(x)$. The set $A(g') = I_{g'}^+(x) \cap I_{g'}^-(x) \cap S \cap I_{\bar{g}}^+(x) \neq \emptyset$ is open in the topology inherited by S and non-empty because $\sigma_{g'}$ must meet $S \cap I_{\bar{g}}^+(x)$. In the topology of S , $\bar{A}(g')$ are compact and $\bigcap_{g < g' < \bar{g}} \bar{A}(g') \neq \emptyset$.

This result follows because otherwise $\bigcup_{g < g' < \bar{g}} \bar{A}^C(g') = S$ where the complement C is taken in the topological space S . Since $\bar{A}^C(g')$ are open sets there would be a finite covering $S = \bar{A}^C(g_1) \cup \dots \cup \bar{A}^C(g_k)$. Now, note that if $\hat{g} < \tilde{g}$ then, since a timelike curve for $\hat{g}$ is timelike for $\tilde{g}$ it is $A(\hat{g}) \subset A(\tilde{g})$ and $\bar{A}^C(\hat{g}) \supset \bar{A}^C(\tilde{g})$. A metric $\check{g}$ exists such that for $i = 1, \dots, k$, $g < \check{g} < g_i$, thus $\bar{A}^C(\check{g}) \supset \bigcup_{i=1}^k \bar{A}^C(g_i) = S$, hence $A(\check{g}) = \emptyset$ a contradiction.

Thus there is $z \in \bigcap_{g < g' < \bar{g}} \bar{A}(g') \neq \emptyset$. In other words there is an event $z \in S$ such that for every $g' > g$ there are (g') -timelike curves starting from x and passing arbitrarily close to z . Thus for every $g' > g$, $(x, z) \in \bar{J}_{g'}^+$ and $(z, x) \in \bar{J}_{g'}^+$ thus by lemma 3.3, $(x, z) \in J_S^+$ and $(z, x) \in J_S^+$ but $x \neq z$, i.e. J_S^+ is not a partial order. $\square$




Figure 1: The idea behind the proof of lemma 3.11. If the event x of the statement does not exist there is a sequence of compacts C_n and metrics $g_n > g$ such that (M, g_n) is chronological in $A_n \supset C_n$. Then a metric $g' > g$ exists which is chronological everywhere (in contradiction with the non-stable causality of (M, g)), indeed, if not, a closed timelike curve γ would have a point $p \in \bar{B}_i$ with i lowest possible index, then (M, g_i) would not be chronological at $p \in A_i$ (in the figure $i = 2$) a contradiction.

Lemma 3.10. *If (M, g) is chronological at x then for every $g' < g$, (M, g') is strongly causal at x . (Stated in another way, if (M, g') is non-strongly causal at x then for every $g > g'$, there is a (g) -timelike closed curve passing through x .)*

Proof. If (M, g') is not strongly causal at x then the characterizing property (ii) of [6, Lemma 3.22] does not hold, that is, there is a neighborhood $U \ni x$ and a sequence of (g') -causal curves σ_n of endpoints x_n, z_n , with $x_n \rightarrow x$, $z_n \rightarrow x$, not entirely contained in U . Let $C \ni x$ be a (g') -convex neighborhood whose compact closure is contained in another (g') -convex neighborhood $V \subset U$ contained in a globally hyperbolic neighborhood. Let $c_n \in \dot{C}$ be the first point at which σ_n escapes C , and let d_n be the last point at which σ_n reenters C . Since $\dot{C}$ is compact there are $c, d \in \dot{C}$, and a subsequence σ_k such that $c_k \rightarrow c$, $d_k \rightarrow d$ and since V is convex, the causal relation on $V \times V$, $J_{(V, g')}^+$, is closed and hence $(x, c), (d, x) \in J_{(V, g')}^+$ thus $(x, c), (d, x) \in J_{g'}^+$ (note that $(d, c) \in J_{(V, g')}^+$ thus d and c must be distinct since the spacetime (V, g') is causal). Taking into account that $(c_k, d_k) \in J_{g'}^+$ it is $(c, d) \in \bar{J}_{g'}^+$. Thus there is a (g) -timelike curve connecting d to c passing through x , and since $(c, d) \in \bar{J}_g^+$ and I_g^+ is open there is a closed timelike curve passing through x . $\square$

Lemma 3.11. *If for every $x \in M$ there is a $(x$ dependent) $g_x > g$ such that (M, g_x) is chronological at x then (M, g) is stably causal. (Stated in another way, if (M, g) is non-stably causal then there is an event $x \in M$ such that for every $\bar{g} > g$, $(M, \bar{g})$ is non-chronological at x .)*

Proof. Let (M, g) be non-stably causal and assume by contradiction that for every $y \in M$ there is a (y dependent) $\bar{g}_y > g$ such that $(M, \bar{g}_y)$ is chronological at y . By Lemma 3.10, taken g_y such that $g < g_y < \bar{g}_y$, (M, g_y) is strongly causal at y and hence it is strongly causal in an open neighborhood U_y of y [8].

Let C be a compact. From the open covering $\{U_y, y \in C\}$, a finite covering can be extracted $\{U_{y_1}, U_{y_2}, \dots, U_{y_k}\}$. A metric $g_C > g$, on M can be found such that for $i = 1, \dots, k$, $g_C < g_{y_i}$ on M . Thus (M, g_C) is strongly causal, and hence chronological, on a open set $A = \cup_i U_{y_i} \supset C$. Let (g_n, C_n, A_n) be a sequence of metrics $g_n > g$, $g_{n+1} < g_n$, and strictly increasing compacts and open sets $C_n \subset A_n \subset C_{n+1}$, such that (M, g_n) is chronological on A_n , and $\cup_n C_n = M$ (for instance introduce a complete Riemannian metric and define C_n as the balls of radius n centered at $x_0 \in M$, $C_n = B(x_0, n)$). Let $\chi_n : M \rightarrow [0, 1]$ be smooth functions such that $\chi_n = 1$ on C_n , and $\chi_n = 0$ outside an open set B_n such that

$$\dots \subset C_n \subset B_n \subset \bar{B}_n \subset A_n \subset C_{n+1} \subset B_{n+1} \subset \bar{B}_{n+1} \subset A_{n+1} \subset \dots$$

Construct a metric $g' > g$ on M as follows (see figure 1). The metric g' on $C_{n+1} \setminus B_n$ has value g_{n+1} , and on $B_n \setminus C_n$ has value $\chi_n g_n + (1 - \chi_n) g_{n+1}$.

The spacetime (M, g') is chronological otherwise there would be a closed (g' -)timelike curve γ . Let i be the minimum integer such that $\bar{B}_i \cap \gamma \neq \emptyset$, and let $p \in \bar{B}_i \cap \gamma$. Then γ is also a closed (g_i -)timelike curve in (M, g_i) , thus chronology is violated at $p \in \bar{B}_i \subset A_i$ a contradiction.

Thus (M, g') is chronological, and hence (M, g) is stably chronological, or equivalently, stably causal (corollary 2.3). $\square$

The next theorem was stated by Seifert [10, Lemma 1]. Unfortunately, he did not give many details and I would say that his argument can not be considered a proof. Hawking and Sachs gave another proof² (see the proof of [4, Prop. 2.3]). The proof given here differs from those and takes advantage of the previous lemmas.

Theorem 3.12. *The relation J_S^+ on $M \times M$ is a partial order if and only if (M, g) is stably causal.*

Proof. It is trivial that if (M, g) is stably causal then J_S^+ is a partial order. Indeed $(x, y) \in J_S^+$ and $(y, x) \in J_S^+$ imply that for a suitable $g' > g$, such that (M, g') is causal, $(x, y) \in J_{g'}^+$ and $(y, x) \in J_{g'}^+$, hence $x = y$ because of the causality of (M, g') .

For the converse let J_S^+ be a partial order, then for every $x \in M$ there is (lemma 3.9) a x -dependent metric $g_x > g$ such that (M, g_x) is chronological at x , thus (M, g) is stably causal because of lemma 3.11.

²There seems to be a gap in Hawking and Sachs's proof. At the very beginning they state that given the spacetime (M, g) and $x \in M$, if J_S^+ is a partial order then there is some $\bar{g} > g$ such that $(M, \bar{g})$ is causal at x . However, they give no argument for this claim. It seems to me that since J_S^+ is a partial order then for every $z \in M$, there is a $\bar{g}_z > g$ such that $(M, \bar{g}_z)$ has no closed causal curve which passes through x and z , but, without a proof of the contrary, $\bar{g}_z$ may well depend on z . Also note that if the claim were obvious then lemma 3.11 would suffice to prove the theorem. This gap is answered by lemma 3.9.

□

From corollary 3.5 and theorem 3.12 it follows

Corollary 3.13. *If (M, g) is stably causal then it is K -causal.*

4 Violating sets on M and $M \times M$

In this section some results on the violating sets for J_S^+ are obtained.

Lemma 4.1. *The stable causality violating set on $M \times M$ for the spacetime (M, g) is the intersection of the chronological violating sets on $M \times M$ for $g' > g$, namely*

$$VJ_{Sg} = \bigcap_{g' > g} VI_{g'}, \quad (2)$$

moreover $VJ_{Sg} = \bigcap_{g' > g} VJ_{g'} = \bigcap_{g' > g} V\bar{J}_{g'} = \bigcap_{g' > g} VK_{g'}$. Finally, VI is open, while $VJ_S \cup \Delta$, $V\bar{J} \cup \Delta$ and $VK \cup \Delta$ are closed.

Proof. From lemma 3.3 $J_{Sg}^+ = \Delta \cup \bigcap_{g' > g} I_{g'}^+$ thus

$$J_{Sg}^+ \cap J_{Sg}^- \cap \Delta^C = \left(\bigcap_{g' > g} I_{g'}^+ \right) \cap \left(\bigcap_{g' > g} I_{g'}^- \right) \cap \Delta^C = \bigcap_{g' > g} (I_{g'}^+ \cap I_{g'}^- \cap \Delta^C).$$

The other equations are proved analogously, the last one using lemma 3.7. It has been already mentioned that since I^+ is open VI is open. Since $VK \cup \Delta = K^+ \cap K^-$ this set is closed and an analogous argument holds for $V\bar{J} \cup \Delta$ and $VJ_S \cup \Delta$. □

The original definition of stable causality implies that if (M, g) is non-stably causal then for every $g' > g$ there is a (g -dependent) event $x_{g'} \in M$ and a (g' -)timelike closed curve through it. Actually, the equivalence between stable causality and the property of antisymmetry for J_S^+ , together with lemma 4.1 imply a considerably stronger result

Corollary 4.2. *If (M, g) is non-stably causal then there is $(x, z) \in M \times M$, $x \neq z$, such that for every $g' > g$, $(x, z) \in I_{g'}^+$ and $(z, x) \in I_{g'}^+$.*

Lemma 4.3. *The stable causality violating set on M for the spacetime (M, g) is the intersection of the chronological violating sets on M for $g' > g$, namely*

$$vJ_{Sg} = \bigcap_{g' > g} vI_{g'}, \quad (3)$$

moreover $vJ_{Sg} = \bigcap_{g' > g} vJ_{g'} = \bigcap_{g' > g} v\bar{J}_{g'} = \bigcap_{g' > g} vK_{g'}$. Finally, vI is open, while vJ_S and vK are closed (the proof of the closure of these sets will follow from lemma 5.9).

Proof. From $vJ_{Sg} = \pi(VJ_{Sg}) = \pi(\bigcap_{g' > g} VI_{g'}) \subset \bigcap_{g' > g} \pi(VI_{g'}) = \bigcap_{g' > g} vI_{g'}$ an inclusion is obtained. The other direction follows by noticing that if $x \in \bigcap_{g' > g} vI_{g'}$ then for every $g' > g$ there is a (g') -timelike closed curve passing through x . The proof of lemma 3.9 shows that under the same assumptions the existence of an event $z \neq x$ can be inferred, such that $(x, z) \in J_{Sg}^+$ and $(z, x) \in J_{Sg}^+$, that is $(x, z) \in VJ_{Sg}$, and finally $x \in vJ_{Sg}$.

Projecting $VI \subset VJ \subset V\bar{J} \subset VK$ we obtain $vI \subset vJ \subset v\bar{J} \subset vK$, which implies $\bigcap_{g' > g} vI_{g'} \subset \bigcap_{g' > g} vJ_{g'} \subset \bigcap_{g' > g} v\bar{J}_{g'} \subset \bigcap_{g' > g} vK_{g'}$. Recall from lemma 3.6 that if $g < g'$ then $J_g^+ \subset \bar{J}_g^+ \subset K_g^+ \subset \Delta \cup I_{g'}^+$, thus $VJ_g \subset V\bar{J}_g \subset VK_g \subset VI_{g'}$ and projecting $vJ_g \subset v\bar{J}_g \subset vK_g \subset vI_{g'}$. We can now prove all the other inclusions. I am going to prove that for every $g' > g$, $\bigcap_{\bar{g} > g} vK_{\bar{g}} \subset vI_{g'}$, indeed I can always find $\tilde{g}$, $g < \tilde{g} < g'$, and from $\bigcap_{\bar{g} > g} vK_{\bar{g}} \subset vK_{\tilde{g}} \subset vI_{g'}$ the thesis follows. Thus since for every $g' > g$, $\bigcap_{\bar{g} > g} vK_{\bar{g}} \subset vI_{g'}$, it follows $\bigcap_{\bar{g} > g} vK_{\bar{g}} \subset \bigcap_{g' > g} vI_{g'}$. The equations are proved.

It is well known that vI is open [8, 4.26], that vK is closed will be proved in lemma 5.9 and from $vJ_{Sg} = \bigcap_{g' > g} vK_{g'}$ it follows the closure of vJ_{Sg} . $\square$

Theorem 4.4. $vI = M$ iff $I^+ = M \times M$, analogously, $vJ_S = M$ iff $J_S^+ = M \times M$.

Proof. The first statement is well known, indeed $vI \neq M$ and $I^+ \neq M \times M$ are two equivalent definitions of a non-totally vicious spacetime. The implication $I^+ = M \times M \Rightarrow vI = M$ is obvious, the other direction follows because vI is made of disjoint open components and for every p, q in a same component $p \ll q$. If $vI = M$ there is only one component which coincides with M .

Let me prove the non-trivial part of the last statement, namely $vJ_S = M \Rightarrow J_S^+ = M \times M$. Indeed, $M = vJ_S = \bigcap_{g' > g} vI_{g'} \Rightarrow \forall g' > g, vI_{g'} = M \Rightarrow I_{g'}^+ = M \times M$ from the first statement and hence $J_S^+ = \bigcap_{g' > g} I_{g'}^+ = M \times M$. $\square$

Remark 4.5. Contrary to non-total viciousness which stays at the bottom of the causal ladder, the condition $vJ_S \neq M$ has no place in the causal hierarchy. Indeed, a causal spacetime may have $vJ_S = M$ (example 5.2 below), which may suggest that perhaps the property $vJ_S \neq M$ is stronger than causality. However, it is easy to give examples of non-chronological non-totally vicious spacetimes with $vJ_S \neq M$ (identify two spacelike parallel lines in Minkowski spacetime to obtain a cylinder and remove from it two parallel spacelike half lines).

A good name for the condition $vJ_S \neq M$ is *stable non-total viciousness*, because it holds iff $\exists g' > g : vI_{g'} \neq M$. Nevertheless, it can also be *non-total non-stable causality*, because $vJ_S \neq \emptyset$ denotes non-stable causality, and $vJ_S \neq M$ states that this non-stable causality is not total.

5 K -causality and stable causality

An important question is whether it is always $K^+ = J_S^+$. The answer is negative as the next examples prove

Example 5.1. Consider the 1+1 cylindrical flat spacetime $M = \mathbb{R} \times S^1$, of metric $ds^2 = -dyd\theta$, $y \in \mathbb{R}$, $\theta \in [0, 2\pi)$. This spacetime is non-causal (hence non-distinguishing) and reflecting, moreover given $x \in M$, $K^+(x) = J^+(x) = \{z \in M : y(z) \geq y(x)\} \neq M$ while $J_S^+(x) = M$ hence $K^+ \neq J_S^+$.

Example 5.2. The 2+1 spacetime $M = \mathbb{R} \times S^1 \times S^1$, of metric $ds^2 = -dyd\theta + d\phi^2$, $\theta, \phi \in \mathbb{R}$, with the identifications $(y, \theta, \phi) = (y, \theta, \phi + 1)$ and $(y, \theta, \phi) = (y, \theta + 1, \phi + \alpha)$, α irrational number is a causal non-distinguishing reflecting spacetime for which given $x \in M$, $K^+(x) = \overline{J^+(x)} = \{z \in M : y(z) \geq y(x)\} \neq M$ while $J_S^+(x) = M$ hence $K^+ \neq J_S^+$. The fact that $J_S^+ = M$ follows for the same reason of the previous example namely the compactness and lightlike nature of the space section which in this case is a torus.

Although K^+ is not always coincident with J_S^+ it can be that K -causality coincides with stable causality. For instance this may happen because when the spacetime is K -causal the two causal relations coincide as stated by claim 1.2.

In order to proceed we have to prove some statements regarding the K^+ relation. First, recall that every event x of the spacetime (M, g) admits an arbitrarily small convex neighborhoods (and arbitrarily small globally hyperbolic neighborhoods). If U is such a neighborhood the causal relation on the spacetime (U, g) , $J_U^+ \subset U \times U$, is closed and hence coincides with the relation K_U^+ . This observation shows that if it were not for global aspects the relation K^+ would be quite simple.

The next two lemmas were proved by Sorkin and Woolgar³ [11, Lemmas 14,15] using a *transfinite induction* argument. Here I give different proofs which use only soft topological methods. I show that in most cases the transfinite induction argument can be avoided. The suggested general strategy is as follows. First, convert the property to be proved into a causal relation on $M \times M$, then show that it is transitive, closed and that contains J^+ , finally use the minimality of K^+ . This approach is particularly clear if the statement of the theorem is rearranged in a suitable way which reads “Let $(x, z) \in K^+$, if *hypothesis* then *thesis*”.

Lemma 5.3. *Let $B \subset M$ be an open set of compact boundary $\dot{B}$. Let $(x, z) \in K^+$, if $x \in B$ and $z \notin B$ (or viceversa), then there is $y \in \dot{B}$ such that $(x, y) \in K^+$ and $(y, z) \in K^+$.*

Proof. Let $T^+ \subset K^+$ be the set of pairs $(x, z) \in K^+$ at which the statement of the theorem is true. This may happen for instance because the hypothesis “ $x \in B$ and $z \notin B$ (or viceversa)” is false or because the hypothesis is true and the thesis “there is $y \in \dot{B}$ such that $(x, y) \in K^+$ and $(y, z) \in K^+$ ” is true.

³The version given here has slightly weaker assumptions because the set B is not required to be compact but only to have compact boundary. This difference will be important in the following.

It is $J^+ \subset T^+$ because if $(x, z) \in J^+$ and the hypothesis “ $x \in B, z \notin B$ (or viceversa)” is true then the thesis is true, y being the intersection of the causal curve σ connecting x to z with $\dot{B}$ (the map $\sigma : [0, 1] \rightarrow M$ is defined over a compact, the set $\sigma^{-1}(\dot{B})$ being closed and limited is a compact and hence there is a last point y at which the curve escapes $\dot{B}$).

Also T^+ is closed, indeed if $(x, z) \in \bar{T}^+$ then either “ $x \in B$ and $z \notin B$ (or viceversa)” is false, in which case $(x, z) \in T^+$ and there is nothing else to prove or “ $x \in B$ and $z \notin B$ (or viceversa)” is true. Assume $x \in B$ and $z \notin B$ the other case being analogous. There is a sequence $(x_k, z_k) \in T^+ \subset K^+$, $(x_k, z_k) \rightarrow (x, z)$, and for sufficiently large k , $x_k \in B$. Now, if $z \in \dot{B}$, then $(x, z) \in T^+$ because it satisfies the thesis of the theorem with $y = z$. Thus we are left with the case $z \in \bar{B}^C$ which is an open set, and hence for sufficiently large k , $z_k \notin B$. Since $(x_k, z_k) \in T^+$, and the hypothesis “ $x_k \in B$ and $z_k \notin B$ (or viceversa)” is satisfied, there are $y_k \in \dot{B}$, $(x_k, y_k) \in K^+$ and $(y_k, z_k) \in K^+$. Then there is an accumulation point $y \in \dot{B}$ and since K^+ is closed, $(x, y) \in K^+$ and $(y, z) \in K^+$ which implies that $(x, z) \in T^+$ because the thesis of the implication is true.

Finally, T^+ is transitive. Indeed, let $(x, w) \in T^+$ and $(w, z) \in T^+$ then the only way in which (x, z) could not belong to T^+ is if $x \in B, z \notin B$ (or viceversa) and the thesis is false. However, in this case w must either belong to B or to B^C , in the former case since $(w, z) \in T^+$ there must be the sought $y \in \dot{B}$, $(x, w) \in K^+$, $(w, y) \in K^+$ and $(y, z) \in K^+$ so that the thesis is verified because $(x, y) \in K^+$. The latter case is analogous.

Thus $T^+ \subset K^+$ is closed, transitive and contains J^+ . By the minimality of K^+ it is $T^+ = K^+$ and hence the implication of the theorem is true for every $(x, z) \in K^+$. □

Let K_B^+ the the K^+ relation for the spacetime (B, g) . It can be regarded not only as a subset of $B \times B$ but also, through the natural inclusion, as a subset of $M \times M$.

Lemma 5.4. *Let $B \subset M$ be an open set of compact boundary $\dot{B}$. Let $(x, z) \in K^+$, if $x, z \in B$ and $(x, z) \notin K_B^+$ then there is $y \in \dot{B}$ such that $(x, y) \in K^+$ and $(y, z) \in K^+$.*

Proof. Let $T^+ \subset K^+$ be the set of pairs $(x, z) \in K^+$ at which the statement of the theorem is true. This may happen for instance because the hypothesis “ $x, z \in B$ and $(x, z) \notin K_B^+$ ” is false or because the hypothesis is true and the thesis “there is $y \in \dot{B}$ such that $(x, y) \in K^+$ and $(y, z) \in K^+$ ” is true.

It is $J^+ \subset T^+$ because if $(x, z) \in J^+$ and the hypothesis “ $x, z \in B$ and $(x, z) \notin K_B^+$ ” is also true then the thesis is true, y being a point in the intersection between the causal curve connecting x to z and $\dot{B}$. The causal curve can not be entirely contained in B otherwise $(x, z) \in J_B^+ \subset K_B^+$.

Also T^+ is closed, indeed if $(x, z) \in \bar{T}^+$ then either “ $x, z \in B$ and $(x, z) \notin K_B^+$ ” is false, in which case $(x, z) \in T^+$ and there is nothing else to prove or “ $x, z \in B$ and $(x, z) \notin K_B^+$ ” is true. Let $x, z \in B$ and $(x, z) \notin K_B^+$ and let

$(x_k, z_k) \in T^+$, be a sequence such that $(x_k, z_k) \rightarrow (x, z)$. Since B is open, for sufficiently large k , $x_k, z_k \in B$, moreover we can assume $(x_k, z_k) \notin K_B^+$. Indeed, if there is a subsequence $(x_s, z_s) \in K_B^+$ then $(x, z) \in K_B^+$ and hence $(x, z) \in T^+$ because the hypothesis is false. Thus $x_k, z_k \in B$ and $(x_k, z_k) \notin K_B^+$ and since $(x_k, z_k) \in T^+$ there are $y_k \in \dot{B}$, such that $(x_k, y_k) \in K^+$ and $(y_k, z_k) \in K^+$. Then there is an accumulation point $y \in \dot{B}$ and since K^+ is closed, $(x, y) \in K^+$ and $(y, z) \in K^+$ which implies that $(x, z) \in T^+$ because the thesis of the implication is true.

Finally, T^+ is transitive. Indeed, let $(x, w) \in T^+$ and $(w, z) \in T^+$ then the only way in which (x, z) could not belong to T^+ is if $x, z \in B$, $(x, z) \notin K_B^+$, and the thesis is false. However, in this case w must either belong to B or to B^C , in the former case if $(x, w) \in K_B^+$ and $(w, z) \in K_B^+$ then $(x, z) \in K_B^+$ and hence $(x, z) \in T^+$. If instead, say $(x, w) \notin K_B^+$ (the other case being analogous), then since $(x, w) \in T^+$ and the thesis is true for (x, w) there is $y \in \dot{B}$, $(x, y) \in K^+$ and $(y, w) \in K^+$ from which $(y, z) \in K^+$ and $(x, z) \in T^+$ follows. If instead $w \in B^C$, then lemma 5.3 can be applied to (x, w) to infer the existence of $y \in \dot{B}$ as required by the thesis.

Thus $T^+ \subset K^+$ is closed, transitive and contains J^+ . By the minimality of K^+ it is $T^+ = K^+$ and hence the implication of the theorem is true for every $(x, z) \in K^+$. □

The next result has been proved by Sorkin and Woolgar [11, Lemma 16] by making use of the previous lemmas and represents the first step in Seifert's proof program. For completeness I include the proof.

Lemma 5.5. *If (M, g) is K -causal at x then x admits arbitrarily small K -convex neighborhoods. In particular, if (M, g) is K -causal then it is strongly K -causal.*

Proof. Given $x \in M$ and $N \ni x$ a arbitrary neighborhood, there is always a strongly causal simple neighborhood $V \subset N$, $x \in V$ (see, for instance, [6, Sect. 2.3]; recall that a simple neighborhood is a convex neighborhood of compact closure contained in another convex neighborhood [8, Sect. 1]). Since V is convex $K_V^+ = J_V^+$. Let $U_n \ni x$, $\bar{U}_{n+1} \subset U_n$, be a sequence of neighborhoods causally convex (and hence K -convex) with respect to V . Let them be a base for the topology at x , namely each open set containing x contains one U_n .

Assume there is a subsequence U_k of non- K -convex neighborhoods. There are $x_k, z_k \in U_k$, and $y_k \notin U_k$, such that $(x_k, y_k) \in K^+$ and $(y_k, z_k) \in K^+$. The event y_k belongs or not to V . In the former case it cannot be $(x_k, y_k) \in K_V^+$ and $(y_k, z_k) \in K_V^+$ because U_k is K_V -convex, thus there is a pair among (x_k, y_k) and (y_k, z_k) to which lemma 5.4 can be applied. The result is the existence of $w_k \in \dot{V}$ such that $(x_k, w_k) \in K^+$ and $(w_k, z_k) \in K^+$. In the latter case the application of lemma 5.3 gives again the existence of $w_k \in \dot{V}$ such that $(x_k, w_k) \in K^+$ and $(w_k, z_k) \in K^+$. Let $w \in \dot{B}$ be an accumulation point of the sequence w_k , then since $x_k, z_k \rightarrow x$, $(x, y) \in K^+$ and $(y, x) \in K^+$ in contradiction with the

K -causality at x . Thus for sufficiently large n all the open sets $U_n \subset N$ must be K -convex neighborhoods. $\square$

Remark 5.6. Note that K -causality implies strong K -causality instead of only the strong $K_{g'}$ -causality for $g' < g$ as one would expect from analogy with lemma 3.10. The reason lies in the fact that K^+ is closed (in the proof of lemma 3.10 we could not infer $(c, d) \in J_{g'}^+$ although $(c, d) \in \bar{J}_{g'}^+$, we had instead to pass to $g > g'$ in order to close the causal chain).

The next nice result is due to Sorkin and Woolgar [11, Lemmas 12,13]. The proof I give shows that one can use the abstract notation $\circ$ for the composition so as to take advantage of the distributive property with respect to unions of sets.

Lemma 5.7. *If U is a open subset of M then $K_U^+ \subset K^+|_{U \times U}$. Moreover, if U is also K -convex then $K_U^+ = K^+|_{U \times U}$.*

Proof. $K^+|_{U \times U}$ is closed (in the topology of $U \times U$), transitive and contains $J^+|_{U \times U}$, in particular, $J_U^+ \subset J^+|_{U \times U}$, thus because of the minimality of K_U^+ , $K_U^+ \subset K^+|_{U \times U}$.

The set K_U^+ can be regarded as a subset of $M \times M$ through the natural inclusion. Consider the causal relation on M

$$\tilde{K}^+ = K_U^+ \cup K^+|_{(U \times U)^c}.$$

First note that $J_U^+ \subset K_U^+$ and $J^+|_{(U \times U)^c} \subset K^+|_{(U \times U)^c}$. Moreover, since K -convexity implies causal convexity $J_U^+ = J^+|_{U \times U}$, thus $J^+ \subset \tilde{K}^+$.

Since $K^+|_{(U \times U)^c} = K^+ \cap (U \times U)^c$ and U is open, this term is a closed set. The first term of $\tilde{K}^+$, i.e. K_U^+ , is closed in the topology of $U \times U$ which is that induced from $M \times M$, thus it is closed in $M \times M$, but, possibly, for accumulation points in $\bar{K}_U^+ \cap (U \times U)^c$, however $\bar{K}_U^+ \subset \bar{K}^+ = K^+$ thus these points belong to the second term and hence $\tilde{K}^+$ is closed.

Now, recall that K_U^+ is transitive, $K_U^+ \circ K_U^+ \subset K_U^+$. The K -convexity of U implies $K^+|_{(U \times U)^c} \circ K^+|_{(U \times U)^c} \subset K^+|_{(U \times U)^c}$. Moreover, $K_U^+ \circ K^+|_{(U \times U)^c} \subset K^+|_{(U \times U)^c}$ because the first endpoint is not in U , and analogously with the factors exchanged. The transitivity property for $\tilde{K}^+$ is proved using the distributivity property of $\circ$

$$\begin{aligned} \tilde{K}^+ \circ \tilde{K}^+ &= [K_U^+ \circ K_U^+] \cup [K_U^+ \circ K^+|_{(U \times U)^c}] \cup [K^+|_{(U \times U)^c} \circ K_U^+] \\ &\quad \cup [K^+|_{(U \times U)^c} \circ K^+|_{(U \times U)^c}] \subset \tilde{K}^+. \end{aligned}$$

Thus $\tilde{K}^+ \subset K^+$, $J^+ \subset \tilde{K}^+$ and $\tilde{K}^+$ is transitive and closed thus, because of the minimality of K^+ , $\tilde{K}^+ = K^+$, and hence $K^+|_{U \times U} = \tilde{K}^+|_{U \times U} = K_U^+$. $\square$

Lemma 5.8. *Let (M, g) be a spacetime, and let U be an open K -convex set of compact closure contained in a convex open set V , then K -causality holds at every point of U .*

Proof. Assume there is $x \in U$ at which K -causality does not hold, then there is $z \in M$, $z \neq x$, such that $(x, z) \in K^+$ and $(z, x) \in K^+$. If $z \notin U$ then since $z \in K^+(x) \cap K^-(x)$, U would not be a K -convex. Thus $z \in U$. Because of lemma 5.7, it must be $(x, z) \in K_U^+$ and $(z, x) \in K_U^+$ but this is impossible because $J_V^+|_{U \times U}$ is closed (in the topology of $U \times U$ as it is already closed in the topology of $V \times V$, the set V being convex [7]), transitive and contains J_U^+ , thus $K_U^+ \subset J_V^+|_{U \times U}$ and hence $(x, z) \in J_V^+$ and $(z, x) \in J_V^+$ in contradiction with the causality of every convex neighborhood (it follows from the fact that in a convex neighborhood every pair of causally related events is connected by a unique geodesic of well defined time orientation, alternatively take V inside a causal neighborhood). $\square$

Theorem 5.9. *The set $(vK)^C \subset M$ at which (M, g) is K -causal is open.*

Proof. Assume (M, g) is K -causal at x , and let $V \ni x$ be a convex set. The set V exists and moreover (V, g) is causal. Let U be a open K -convex set, $x \in U$, of compact closure contained in V . It exists because of lemma 5.5. Thus by lemma 5.8 K -causality holds at every point of U and hence $(vK)^C$ is open. $\square$

Lemma 5.10. *If $g_1 \leq g_2$ then $K_{g_1}^+ \subset K_{g_2}^+$. In particular, if K_{g_2} -causality holds at $x \in M$ then K_{g_1} -causality holds at x .*

Proof. $K_{g_2}^+ \subset M \times M$ is (a) closed, (b) transitive and contains $J_{g_2}^+ \supset J_{g_1}^+$ and hence, (c) contains $J_{g_1}^+$, thus must be larger than the smallest set which satisfies (a), (b) and (c), namely $K_{g_1}^+$. $\square$

Lemma 5.11. *Let $B \subset M$ be an open set of compact closure. Let $N = M \setminus \bar{B}$, if $x, z \in N$, $(x, z) \in K^+$ and $(x, z) \notin K_N^+$ then there is $y \in \dot{B}$ such that $(x, y) \in K^+$ and $(y, z) \in K^+$.*

Proof. It is an immediate consequence of lemma 5.4, indeed N is open, is such that $M \setminus N$ is compact and plays the role of the set B in the statement of lemma 5.4 (note that $\dot{N} = \dot{B}$). $\square$

Lemma 5.12. *Let $B \subset M$ be an open set of compact closure. Let $\bar{g} \geq g$, with $\{w \in M : \bar{g}(w) \neq g(w)\} \subset B$, let K_g -causality hold at x , $x \notin \bar{B}$, then if $K_{\bar{g}}$ -causality does not hold at x there is $z \in \bar{B}$ at which $K_{\bar{g}}$ -causality fails too.*

Proof. Since $K_{\bar{g}}$ -causality does not hold at x there must be $z \in M$, $z \neq x$, such that $(x, z) \in K_{\bar{g}}^+$ and $(z, x) \in K_{\bar{g}}^+$. Assume $z \notin \bar{B}$ otherwise there is nothing to prove. Since K_g -causality holds at x , $(x, z) \notin K_g^+$ or $(z, x) \notin K_g^+$. Consider the former case, the other being analogous. Let $N = M \setminus \bar{B}$, from lemma 5.7 it is $K_g^+|_{N \times N} \supset K_{(N, g)}^+ = K_{(N, \bar{g})}^+$ so that $(x, z) \notin K_{(N, \bar{g})}^+$. But it is also $(x, z) \in K_{\bar{g}}^+$ and $x, z \in N$, thus by lemma 5.11 there is $y \in \dot{B}$ such that $(x, y) \in K_{\bar{g}}^+$ and $(y, z) \in K_{\bar{g}}^+$. Composing these relations with $(z, x) \in K_{\bar{g}}^+$, it follows $(x, y) \in K_{\bar{g}}^+$ and $(y, x) \in K_{\bar{g}}^+$ thus $K_{\bar{g}}$ -causality does not hold at $y \in \bar{B}$. $\square$

A fundamental observation is that given a causal relation $R^+ \subset M \times M$, the R -convexity of a set A must be understood as a condition on the shape of the metric on A^C rather than on A . For instance, for J -convexity, the fact that no causal curve can escape and reenter a causally convex set A is a constraint due to the shape of the light cones outside A . The light cone structure inside A has not very much to do with this property. This observation is important because by enlarging the light cones inside a K -convex set one expects to keep the K -causality property. Since a special feature of K -causality is that K -causality implies strong K -causality, the same enlargement can be continued in other places so as to obtain, one would say, a global widening of the light cones. This is basically Seifert's program outlined by him in [10, Lemma 1, point (4)] (note that in that lemma "at least once" is probably a misprint and must be replaced with "at most once"). Unfortunately, in order to follow this program, several technical lemmas are needed. Some have been already proved. The next one is particular because the lengthy proof works only if statements (a1), (a2) and (b) are proved all at the same time.

Lemma 5.13. *Let (M, g) be a spacetime, F a closed set, B an open set of compact boundary $\dot{B}$ and $F \subset B$. Let $(x, z) \in K^+$*

- (a1) *If $x \in \bar{B}$ and $z \notin \bar{B}$ then there is $y \in \dot{B}$ such that $(x, y) \in K^+$ and $(y, z) \in K_{M \setminus F}^+$.*
- (a2) *If $x \notin \bar{B}$ and $z \in \bar{B}$ then there is $y \in \dot{B}$ such that $(x, y) \in K_{M \setminus F}^+$ and $(y, z) \in K^+$.*
- (b) *If $x, z \notin \bar{B}$ then either $(x, z) \in K_{M \setminus F}^+$ or there are $y_1, y_2 \in \dot{B}$ such that $(x, y_1) \in K_{M \setminus F}^+$, $(y_2, z) \in K_{M \setminus F}^+$, and $(y_1, y_2) \in K^+$.*

In particular if $U, \bar{B} \subset U$, is an open set such that $U \setminus F$ is $K_{M \setminus F}$ -convex then U is K -convex.

Proof. Recall that $K_{M \setminus F}^+ \subset K^+$. Let $T^+ \subset K^+$ be the set of pairs $(x, z) \in K^+$ at which all the three statements of the theorem are true (which, selected a statement, may happen because the hypothesis is false or because the thesis is true).

It is $J^+ \subset T^+$ because the statements (a1), (a2) and (b) are all true in this case. Indeed, consider (a1). If $(x, z) \in J^+$ and the hypothesis " $x \in \bar{B}$, $z \notin \bar{B}$ " is also true then the thesis is true, y being the last point of the causal curve connecting x to z in $\bar{B}$ (the segment of the causal curve connecting y to z is entirely contained in $M \setminus F$). The statement (a2) is proved similarly. As for (b), if $(x, z) \in J^+$ and $x, z \notin \bar{B}$ then if the causal curve connecting x to z does not intersect $\bar{B}$ then $(x, z) \in J_{M \setminus F}^+$, otherwise there is a first point y_1 at which the causal curve enters $\bar{B}$ and a last point y_2 at which it leaves $\bar{B}$ so that the segments of causal curves connecting x to y_1 and y_2 to z are contained on $M \setminus F$ and hence $(x, y_1) \in J_{M \setminus F}^+$ and $(y_2, z) \in J_{M \setminus F}^+$ while $(y_1, y_2) \in J^+ \subset K^+$ is obvious.

Also T^+ is closed, indeed if $(x, z) \in \bar{T}^+$ then either $x, z \in \bar{B}$ in which case all the hypothesis of (a1), (a2) and (b) are false, and thus the statements are true, $(x, z) \in T^+$ and there is nothing left to prove, or only one of those mutually excluding hypothesis is true.

Suppose the hypothesis of (a1) is true, that is, $x \in \bar{B}$, $z \notin \bar{B}$. In this case statements (a2) and (b) are true because their hypothesis are false and we have only to check that statement (a1) is true. There is a sequence $(x_k, z_k) \in T^+$, $(x_k, z_k) \rightarrow (x, z)$, and, since $\bar{B}^C$ is open, for sufficiently large k , $z_k \notin \bar{B}$. Now, without loss of generality we can assume (pass to a subsequence if necessary) that either (i) $x_k \in \bar{B}$ or (ii) $x_k \notin \bar{B}$. In case (i) since $(x_k, z_k) \in T^+$, and the hypothesis of (a1) “ $x_k \in \bar{B}$ and $z_k \notin \bar{B}$ ” is satisfied, there are $y_k \in \bar{B}$, $(x_k, y_k) \in K^+$ and $(y_k, z_k) \in K_{M \setminus F}^+$. Then there is an accumulation point $y \in \bar{B}$ and since K^+ and $K_{M \setminus F}^+$ are both closed, $(x, y) \in K^+$ and $(y, z) \in K_{M \setminus F}^+$ which implies that $(x, z) \in T^+$ because the thesis of (a1) and hence statement (a1) is true. In case (ii) since $(x_k, z_k) \in T^+$, and the hypothesis of (b) “ $x_k, z_k \notin \bar{B}$ ” is satisfied either there is a subsequence denoted in the same way such that $(x_k, z_k) \in K_{M \setminus F}^+$ in which case $(x, z) \in K_{M \setminus F}^+$ and the thesis of (a1) is verified with $y = x$, or there are $y_{1k}, y_{2k} \in \bar{B}$, such that $(x_k, y_{1k}) \in K_{M \setminus F}^+$, $(y_{2k}, z_k) \in K_{M \setminus F}^+$ and $(y_{1k}, y_{2k}) \in K^+$. Then there are accumulation points $y_1, y_2 \in \bar{B}$ and since $K_{M \setminus F}^+$ and K^+ are closed, $(x, y_1) \in K_{M \setminus F}^+$, $(y_2, z) \in K_{M \setminus F}^+$ and $(y_1, y_2) \in K^+$. Thus $(x, y_2) \in K^+$ and $(y_2, z) \in K_{M \setminus F}^+$ which implies that $(x, z) \in T^+$ because the thesis of (a1) and hence statement (a1) is true along with (a2) and (b).

The proof assuming true the hypothesis of (a2) is analogous.

Suppose the hypothesis of (b) is true, that is, $x, z \notin \bar{B}$. In this case statements (a1) and (a2) are true because their hypothesis is false and we have only to check that statement (b) is true. There is a sequence $(x_k, z_k) \in T^+$, $(x_k, z_k) \rightarrow (x, z)$, and, since $\bar{B}^C$ is open, for sufficiently large k , $x_k, z_k \notin \bar{B}$. Since $(x_k, z_k) \in T^+$ and the hypothesis of (b) is true, for each (sufficiently large) k either $(x_k, z_k) \in K_{M \setminus F}^+$ or there are $y_{1k}, y_{2k} \in \bar{B}$ such that $(x_k, y_{1k}) \in K_{M \setminus F}^+$, $(y_{2k}, z_k) \in K_{M \setminus F}^+$ and $(y_{1k}, y_{2k}) \in K^+$. If there is a subsequence such that the first possibility holds then $(x, z) \in K_{M \setminus F}^+$ and the thesis of (b) and hence statement (b) is true. Otherwise for all but a finite number of values of k the second possibility holds then there are accumulation points y_1 of y_{1k} and y_2 of y_{2k} so that, because of the closure of $K_{M \setminus F}^+$ and K^+ , $(x, y_1) \in K_{M \setminus F}^+$, $(y_2, z) \in K_{M \setminus F}^+$ and $(y_1, y_2) \in K^+$. Thus, again, (b) is true because its thesis is true.

Finally, T^+ is transitive. Indeed, let $(x, w) \in T^+$ and $(w, z) \in T^+$ then the only way in which (x, z) could not belong to T^+ is if (x, z) contradicts one of the statements (a1), (a2) or (b). Assume this happens for (a1) then $x \in \bar{B}$ and $z \notin \bar{B}$ while the thesis of (a1) is false (note that (a2) and (b) are true because their hypothesis is false). However, in this case w must either belong to $\bar{B}$ or to $\bar{B}^C$, in the former case since $(w, z) \in T^+$ and $w \in \bar{B}$, $z \notin \bar{B}$, there must be $y \in \bar{B}$, $(x, w) \in K^+$, $(w, y) \in K^+$ and $(y, z) \in K_{M \setminus F}^+$ so that the thesis of (a1) is verified because through composition $(x, y) \in K^+$. In the latter case

since $(x, w) \in T^+$ and $x \in \bar{B}$, $w \notin \bar{B}$, there is $y \in \dot{B}$ such that $(x, y) \in K^+$ and $(y, w) \in K_{M \setminus F}^+$. Unfortunately, the composition with $(w, z) \in K^+$ can not be immediately done, however, since $(w, z) \in T^+$ and $w, z \notin \bar{B}$ either $(w, z) \in K_{M \setminus F}^+$, and we have finished because $(x, y) \in K^+$ and $(y, z) \in K_{M \setminus F}^+$ which is the thesis of (a1), or there is $y_2 \in \dot{B}$ (and an analogous y_1 of no interest here) such that $(w, y_2) \in K^+$ and $(y_2, z) \in K_{M \setminus F}^+$. Thus in this last case $(x, y_2) \in K^+$ and $(y_2, z) \in K_{M \setminus F}^+$ that is, the thesis of (a1) is verified. Thus statement (a1) can not be contradicted. The proof for (a2) is analogous.

It remains to show that (b) can not be contradicted by (x, z) . Assume $x, z \notin \bar{B}$, w must either belong to $\bar{B}$ or to $\bar{B}^C$ (note that (a1) and (a2) are true because their hypothesis is false). In the former case since $(x, w) \in T^+$ and $x \notin \bar{B}$, $w \in \bar{B}$, by (a2) there is $y_1 \in \dot{B}$ such that $(x, y_1) \in K_{M \setminus F}^+$, $(y_1, w) \in K^+$. Also, since $(w, z) \in T^+$ and $w \in \bar{B}$, $z \notin \bar{B}$ there is by (a1), $y_2 \in \dot{B}$ such that $(w, y_2) \in K^+$ and $(y_2, z) \in K_{M \setminus F}^+$ thus the thesis of (b) is true for (x, z) as $(y_1, y_2) \in K^+$. It remains to consider the case $w \notin \bar{B}$. In this case $(x, w) \in T^+$, $(w, z) \in T^+$ and $x, w, z \notin \bar{B}$. There are four possibilities depending on which of the cases given by the thesis of (b) applies to the pairs (x, w) and (w, z) . For instance, by (b) it can be $(x, w) \in K_{M \setminus F}^+$. If this is the case also for (w, z) then $(x, z) \in K_{M \setminus F}^+$ and the thesis of (b) holds for (x, z) as required. Otherwise, the second pair may satisfy the second alternative given by (b) namely there are $y_1, y_2 \in \dot{B}$ such that $(w, y_1) \in K_{M \setminus F}^+$, $(y_2, z) \in K_{M \setminus F}^+$ and $(y_1, y_2) \in K^+$. Thus $(x, y_1) \in K_{M \setminus F}^+$, $(y_2, z) \in K_{M \setminus F}^+$ and $(y_1, y_2) \in K^+$ which again makes the thesis of (b) true for (x, z) . There are two cases left.

The case $(x, w) \notin K_{M \setminus F}^+$, $(w, z) \in K_{M \setminus F}^+$ is completely analogous to the one just considered and leads again to the truth of (b). The last case $(x, w) \notin K_{M \setminus F}^+$, $(w, z) \notin K_{M \setminus F}^+$ implies the existence of events $y_1, y_2, \bar{y}_1, \bar{y}_2 \in \dot{B}$ such that $(x, y_1) \in K_{M \setminus F}^+$, $(y_1, y_2) \in K^+$, $(y_2, w) \in K_{M \setminus F}^+$, $(w, \bar{y}_1) \in K_{M \setminus F}^+$, $(\bar{y}_1, \bar{y}_2) \in K^+$ and $(\bar{y}_2, z) \in K_{M \setminus F}^+$. Thus setting $\tilde{y}_1 = y_1$, $\tilde{y}_2 = \bar{y}_2$, it is $(x, \tilde{y}_1) \in K_{M \setminus F}^+$, $(\tilde{y}_1, \tilde{y}_2) \in K^+$ and $(\tilde{y}_2, z) \in K_{M \setminus F}^+$, that is (b) is true for (x, z) . Thus in every case (x, z) can not contradict (b). The transitivity of T^+ is proved.

Thus $T^+ \subset K^+$ is closed, transitive and contains J^+ . By the minimality of K^+ it is $T^+ = K^+$ and hence the implications (a1), (a2) and (b) of the theorem are true for every $(x, z) \in K^+$.

For the last statement of the theorem, if U is not K -convex there are $x, z \in U$, $y \notin U$ (thus $y \notin \bar{B}$) such that $(x, y) \in K^+$ and $(y, z) \in K^+$. Thanks to (a1), (a2) and (b) we can assume without loss of generality, through a redefinition of x and z , $(x, y) \in K_{M \setminus F}^+$ and $(y, z) \in K_{M \setminus F}^+$ which contradicts the $K_{M \setminus F}$ -convexity of $U \setminus F$. $\square$

Lemma 5.14. *Let $F \subset M$ be closed set of compact boundary $\dot{F}$, and $U \supset F$ an open set, then there is an open set B of compact boundary such that $F \subset B \subset \bar{B} \subset U$.*

Proof. For every point $x \in \dot{F}$ it is possible to find a neighborhood $A(x) \ni x$ of compact closure such that $\bar{A}(x) \subset U$. Since $\dot{F}$ is compact there is a finite number of points $x_i, i = 1, \dots, n$, such that $\{A(x_1), \dots, A(x_n)\}$ covers $\dot{F}$. Then the open set $B = F \cup \bigcup_i A(x_i)$ has compact boundary because $\dot{B}$ is a closed subset of the compact $\bigcup_i \bar{A}(x_i)$. Finally, by construction, $F \subset B \subset \bar{B} \subset U$. $\square$

Theorem 5.15. *Let U be an open subset of M and let $F \subset U$ be a closed set. If U is K -convex then $U \setminus F$ is $K_{M \setminus F}$ -convex. Moreover, if F has compact boundary then the converse holds, i.e. if $U \setminus F$ is $K_{M \setminus F}$ -convex then U is K -convex.*

Proof. Regard $K_{M \setminus F}^+$ as a subset of $M \times M$ through the natural inclusion of $(M \setminus F) \times (M \setminus F)$ into $M \times M$. Assume U is K -convex. From lemma 5.7, $K_{M \setminus F}^+ \subset K^+|_{(M \setminus F) \times (M \setminus F)}$. Thus $U \setminus F$ is $K_{M \setminus F}$ -convex. For the converse, by lemma 5.14 there is a open set B of compact boundary $\dot{B}$, such that $F \subset B \subset \bar{B} \subset U$. The conclusion comes from lemma 5.13. $\square$

Theorem 5.16. *Let (M, g) be a spacetime, F a closed set of compact boundary, and $U \supset F$ a open K -convex set. The spacetime is K -causal iff U is K_U -causal and $M \setminus F$ is $K_{M \setminus F}$ -causal.*

Proof. Assume (M, g) is K -causal then since with the usual natural inclusion in $M \times M$, $K_U^+ \subset K^+$, (lemma 5.7) U is K_U -causal and analogously since $K_{M \setminus F}^+ \subset K^+$, $M \setminus F$ is $K_{M \setminus F}$ -causal.

For the converse, assume M is not K -causal then there are $x, z \in M$, $x \neq z$, such that $(x, z) \in K^+$ and $(z, x) \in K^+$. If both are in U then because of lemma 5.7 $(x, z) \in K_U^+$ and $(z, x) \in K_U^+$ which is not possible since U is K_U -causal by assumption. The case $x \in U$, $z \notin U$ implies $z \in K^+(x, x)$ which contradicts the K -convexity of U . The case $x \notin U$, $z \in U$ is analogous.

It remains the case $x, z \notin U$. By lemma 5.14 there is a open set B of compact boundary $\dot{B}$, such that $F \subset B \subset \bar{B} \subset U$, thus $x, z \notin \bar{B}$. Consider the pair $(x, z) \in K^+$ and apply lemma 5.13. It follows that either (i) $(x, z) \in K_{M \setminus F}^+$ or (ii) there are $y_1, y_2 \in U \setminus F$ such that $(x, y_1) \in K_{M \setminus F}^+$, $(y_1, y_2) \in K^+$ and $(y_2, z) \in K_{M \setminus F}^+$. Consider case (ii) and $(z, x) \in K^+$. Applying again lemma 5.13 (iia) $(z, x) \in K_{M \setminus F}^+$ or (iib) there are $\bar{y}_1, \bar{y}_2 \in U \setminus F$ such that $(z, \bar{y}_1) \in K_{M \setminus F}^+$, $(\bar{y}_1, \bar{y}_2) \in K^+$ and $(\bar{y}_2, x) \in K_{M \setminus F}^+$. But case (iia) leads to $(y_2, z) \in K_{M \setminus F}^+$ and $(z, y_1) \in K_{M \setminus F}^+$ in contradiction with the $K_{M \setminus F}$ -convexity of $U \setminus F$. Analogously, case (iib) leads to $(\bar{y}_2, x) \in K_{M \setminus F}^+$ and $(x, y_1) \in K_{M \setminus F}^+$ which again contradicts the $K_{M \setminus F}$ -convexity of $U \setminus F$. Thus $(x, z) \in K_{M \setminus F}^+$ and an analogous reasoning leads to $(z, x) \in K_{M \setminus F}^+$ in contradiction with the $K_{M \setminus F}$ -causality of $M \setminus F$. The overall contradiction proves that M is K -causal. $\square$

Theorem 5.17. *Let (M_1, g_1) and (M_2, g_2) be two spacetimes and let $F_1 \subset M_1$ and $F_2 \subset M_2$ be two closed sets of compact boundaries in the respective*

topologies. Define $N_1 = M_1 \setminus \text{Int } F_1$, $N_2 = M_2 \setminus \text{Int } F_2$, and assume there is an invertible isometry $\phi : N_1 \rightarrow N_2$ of the spacetimes with boundaries $(N_1, g_1|_{N_1})$ and $(N_2, g_2|_{N_2})$, $\phi(F_1) = F_2$. Then open set $U_1 \supset F_1$ is $K_{(M_1, g_1)}$ -convex iff $U_2 = \phi(U_1 \setminus F_1) \cup F_2$ is $K_{(M_2, g_2)}$ -convex.

Assume that U_1 is indeed $K_{(M_1, g_1)}$ -convex, and assume, moreover, that U_1 is $K_{(U_1, g_1|_{U_1})}$ -causal and U_2 is $K_{(U_2, g_2|_{U_2})}$ -causal. Then (M_1, g_1) is $K_{(M_1, g_1)}$ -causal iff (M_2, g_2) is $K_{(M_2, g_2)}$ -causal.

Proof. The set U_1 is $K_{(M_1, g_1)}$ -convex iff $U_1 \setminus F_1$ is $K_{(M_1 \setminus F_1, g_1|_{M_1 \setminus F_1})}$ -convex (theorem 5.15). The map $\phi|_{N_1 \setminus F_1} : M_1 \setminus F_1 \rightarrow M_2 \setminus F_2$ is an isometry thus $U_1 \setminus F_1$ is $K_{(M_1 \setminus F_1, g_1|_{M_1 \setminus F_1})}$ -convex iff $U_2 \setminus F_2 = \phi(U_1 \setminus F_1)$ is $K_{(M_2 \setminus F_2, g_2|_{M_2 \setminus F_2})}$ -convex iff U_2 is $K_{(M_2, g_2)}$ -convex (theorem 5.15).

The last statement follows from theorem 5.16 because, given the assumptions, (M_1, g_1) is $K_{(M_1, g_1)}$ -causal iff $(M_1 \setminus F_1, g_1|_{M_1 \setminus F_1})$ is $K_{(M_1 \setminus F_1, g_1|_{M_1 \setminus F_1})}^+$ -causal which, due to the isometry, holds iff $(M_2 \setminus F_2, g_2|_{M_2 \setminus F_2})$ is $K_{(M_2 \setminus F_2, g_2|_{M_2 \setminus F_2})}^+$ -causal which holds, again by theorem 5.16, iff (M_2, g_2) is $K_{(M_2, g_2)}$ -causal. $\square$

Remark 5.18. The theorem 5.17 is a powerful result which allows to modify the metric and even make surgery operations inside U_1 . In short it states that if a spacetime (M_1, g_1) has a K -causal K -convex open set U_1 , given a closed set of compact boundary $F_1 \subset U_1$ it is possible to arbitrary modify the metric and even the topology inside $\text{Int } F_1$ without altering the K -convexity of “ U_1 ” in the sense that the obtained spacetime (M_2, g_2) will be such that (denoting with C_1 the complement in M_1 and with C_2 the complement in M_2) $U_2 = ((U_1)^{C_1})^{C_2}$ is K -convex.

Even more it states that whatever the metric deformation or surgery operation done inside U_1 , the spacetime does not lose its K -causality provided the attached set F_2 does not make the spacetime $(U_2, g_2|_{U_2})$ non- K -causal.

In the following we shall need to consider metric deformations (theorem 5.19) and even topological surgery operations (theorem 6.2).

Theorem 5.19. *Let (M, g) be a K_g -causal spacetime and let $x \in M$. Then a metric $\bar{g} \geq g$, such that $\bar{g} > g$ in an open neighborhood of x , exists such that the spacetime $(M, \bar{g})$ is $K_{\bar{g}}$ -causal.*

Proof. Since (M, g) is K -causal it is also K -strongly causal. Thus it is always possible to find a open K -convex set $U \ni x$ and a closed set of compact boundary F , $x \in \text{Int } F$, such that $F \subset U$. Even more U can be chosen inside a globally hyperbolic neighborhood V . Since global hyperbolicity implies stable causality and hence stable K -causality the metric can be widened inside F without spoiling the K -causality of U (otherwise the K -causality of V would be spoiled which would be in contradiction with its stable K -causality). Finally, because of theorem 5.17 the resulting spacetime is still K -causal. $\square$

Remark 5.20. The previous result proves that the cones can be widened in a neighborhood of any chosen point without spoiling K -causality. Moreover, the

only assumption of the theorem is K -causality itself thus the procedure can be continued. The cones can be widened in a finite number of points without spoiling K -causality. The problem is that taken a point y each of these enlargements may decrease the size of the K -convex neighborhood around y . In particular the kind of enlargement that can be applied at a neighborhood of a new chosen point could depend on the sequence of enlargements followed previously. Thus, though it would be natural to apply a transfinite induction argument to assure the stable K -causality of (M, g) , an argument in this direction would have to circumvent these technical difficulties.

6 Some consequences of the possible coincidence

Though the proof of the coincidence between K -causality and stable causality has not been given, it is interesting to explore its possible consequences. In this section I will clearly state if this assumption is necessary for the results considered.

Lemma 6.1. *Let (M, g) be a spacetime. If U_1 and U_2 are K -convex sets and $(U_1 \times U_2) \cap (K^+ \cup K^-) = \emptyset$ then $U = U_1 \cup U_2$ is a K -convex set.*

Proof. If U is not K -convex there are $x, z \in U$ and $y \notin U$ such that $(x, y) \in K^+$ and $(y, z) \in K^+$. Now, $x, z \in U_1$ is excluded because U_1 is K -convex, analogously $x, z \in U_2$ is excluded because U_2 is K -convex. Note that $(x, z) \in K^+$ (i.e. $(z, x) \in K^-$). The case $x \in U_1, z \in U_2$, is excluded because $(U_1 \times U_2) \cap K^+ = \emptyset$ and analogously the case $z \in U_1, x \in U_2$, is excluded because $(U_1 \times U_2) \cap K^- = \emptyset$. Thus U is K -convex. $\square$

The next result comes from the development of an idea by H. Seifert [10, Lemma 2].

Theorem 6.2. *Provided K -causality coincides with stable causality: if (M, g) is K -causal (stably causal) then $K^+ = J_S^+$.*

Proof. Assume by contradiction $K_g^+ \neq J_{S_g}^+$ then there is $(x, z) \in J_{S_g}^+ \setminus K_g^+$, and in particular $x \neq z$. It must also be $(z, x) \notin K_g^+$ otherwise $(z, x) \in K_g^+ \subset J_{S_g}^+$, thus $J_{S_g}^+$ would not be a partial order and hence (M, g) would not be stably causal, a contradiction because K -causality and stable causality coincide.

We are going to construct a spacetime which is K -causal and yet non-stably-causal in contradiction with the assumption.

Since $K_g^+ \cup K_g^-$ is closed and $(x, z) \notin K_g^+ \cup K_g^-$ there are open sets of compact closure $V_x \ni x$, and $V_z \ni z$ such that $(V_x \times V_z) \cap (K_g^+ \cup K_g^-) = \emptyset$.

Since (M, g) is K_g -causal there is $\bar{g} > g$ such that $(M, \bar{g})$ is $K_{\bar{g}}$ -causal (equivalence between K -causality, stable causality and stable K -causality, see corollary 2.3) and hence strongly $K_{\bar{g}}$ -causal. Let $U_x \ni x$ (resp. $U_z \ni z$) be a $K_{\bar{g}}$ -convex

neighborhood contained in a $(\bar{g})$ -globally hyperbolic neighborhood W_x (resp. W_z) contained in V_x (resp. V_z). Let u_x be a timelike vector at x , set

$$D_x = \{y \in M : y = \exp_x v, v \in TM_x, g(v, u_x) = 0, 0 < g(v, v) < \epsilon\}$$

where $\epsilon > 0$ is chosen so that the closed disk (ball) $\bar{D}_x$ is contained in U_x , and no $(\bar{g})$ -causal curve starting or ending at x and contained in W_x can intersect $\bar{D}_x$ (this is possible because W_x being globally hyperbolic it is also distinguishing). Let D_z be defined analogously. Let $N = M \setminus (\bar{D}_x \cup \bar{D}_z)$, then $(N, \bar{g}|_N)$ is $K_{(N, \bar{g}|_N)}^+$ -causal as $K_{(N, \bar{g}|_N)}^+ \subset K_{(M, \bar{g})}^+$ (lemma 5.7). Cut M at $\bar{D}_x$, that is let the boundary of N be there topologically split into the two sides D_x^p (above) and D_x^f (below) (and analogously for $\bar{D}_z$).

Consider a globally hyperbolic spacetime (G, g_2) with the topology of $B^{n-1} \times (0, 1)$ where B^{n-1} is the open ball of $\mathbb{R}^{n-1}$ (n is the spacetime dimension). The metric on it can be chosen so that its past boundary at 0 can be glued with D_z^f and its future boundary at 1 can be glued with D_x^p all that preserving the continuity properties of the metric. Call $\tilde{M}$ the final manifold and $\tilde{g}$ be the total metric obtained joining $g|_N$ and g_2 . The actual shape of the metric g_2 is not important, but it must be chosen sufficiently wide so that for every $z' \in D_z^f$, $D_x^p \subset J_{(\tilde{M}, \tilde{g})}^+(z')$.

I claim that the spacetime $(\tilde{M}, \tilde{g})$ is non-stably-causal. Indeed, we know that on M for every $g', g < g' < \bar{g}$, $J_{S_g}^+ \subset \Delta \cup I_{g'}^+$ (lemma 3.3) hence there is a (g') -timelike curve connecting x to z . This curve can not intersect D_x before escaping W_x , nor can it intersect D_z after entering W_z (saved for the endpoints) because of the distinction of W_x, W_z , and because of the very definition of D_x and D_z . However, it can not either intersect D_x after escaping W_x because that would violate the $K_{\bar{g}}$ -convexity of U_x , $D_x \subset U_x \subset W_z$, and analogously for D_z . Every metric $\tilde{g}' > \tilde{g}$ on $\tilde{M}$ induces a metric $\tilde{g}'|_N > g|_N$ on N which can be modified in a neighborhood of D_x and D_z without changing the causal nature of the curves connecting x to z so as to obtain an extension to a metric g' of M . But since any metric g' of this form and such that $g' < \bar{g}$ allows for a (g') -timelike curve from x to z , the same is true for metrics g' which do not satisfy the constraint $g' < \bar{g}$. The conclusion is that for every $\tilde{g}' > \tilde{g}$ there is a (g') -causal and hence $(\tilde{g}')$ -causal curve connecting x to z , and since z and x are also connected by a $\tilde{g}$ -causal curve passing through G , the spacetime $(\tilde{M}, \tilde{g})$ is not stably causal.

It remains to prove that $(\tilde{M}, \tilde{g})$ is K -causal.

By lemma 6.1 $U = U_x \cup U_z$ is $K_{(M, g)}$ -convex and moreover, since (M, g) is K -causal it is $K_{(U, g|_U)}$ -causal (theorem 5.16). The idea is to show that $(\tilde{M}, \tilde{g})$ is obtained from (M, g) through a surgery operation allowed by theorem 5.17 which preserves K -causality. Indeed, it is obvious that it is always possible to find F , closed set of compact boundary, such that $(D_x \cup D_z) \subset F \subset U$. Then in this surgery operation $\text{Int}(F)$ is excised (note that it is disconnected) and replaced by a new set $\text{Int}(\tilde{F})$ (note that it is connected) which is glued in place of it (all this made rigorous by the obvious presence of the isometry cited in theorem 5.17). Intuitively, identifying some sets from M and $\tilde{M}$, $\text{Int}(\tilde{F})$ is the set $\text{Int}(\tilde{F}) =$




Figure 2: The idea behind the proof of theorem 6.2. The dotted disks D_z^p and D_x^f are boundaries of the total spacetime $\tilde{M} = (M \setminus (\bar{D}_x \cup \bar{D}_z)) \cup D_z^f \cup G \cup D_x^p$.

$D_z^f \cup G \cup D_x^p \cup (\text{Int}(F) \setminus (\bar{D}_x \cup \bar{D}_z))$ while $\tilde{U} = D_z^f \cup G \cup D_x^p \cup (U \setminus (\bar{D}_x \cup \bar{D}_z))$. In the same sloppy notation the boundaries of F and $\tilde{F}$ can be identified. Actually, note that because of the inclusion of G , while F was compact, the resulting $\tilde{F}$ is only closed but still of compact boundary which is what is needed for applying theorem 5.17. By theorem 5.17, $\tilde{U}$ is $K_{(\tilde{M}, \tilde{g})}$ -convex. Moreover, $\tilde{U}$ is $K_{(\tilde{U}, \tilde{g}|_{\tilde{U}})}$ -causal because G and the metric on it can be chosen particularly well behaved. In the end all the conditions of theorem 5.17 are met and thus $(\tilde{M}, \tilde{g})$ is $K_{(\tilde{M}, \tilde{g})}$ -causal. $\square$

A spacetime is reflecting if $(x, z) \in \bar{J}^+$ implies $z \in \bar{J}^+(x)$ and $x \in \bar{J}^-(z)$ (this property can be taken as a definition, see [6, Prop. 3.45]). Another equivalent property is $z \in \bar{J}^+(x) \Leftrightarrow x \in \bar{J}^-(z)$.

Recall that a spacetime is *causally continuous* if it is reflecting and distinguishing.

If K -causality coincides with stable causality the next proof is particularly simple. Compare for instance with the original one given by Hawking and Sachs [4]. Actually, it also shows that the full assumptions underlying the definition of causal continuity are not all needed for guaranteeing stable causality.

Theorem 6.3. *Assume that K -causality coincides with stable causality. A spacetime which is future distinguishing and future reflecting (or past distinguishing and past reflecting) is stably causal. In particular causal continuity implies stable causality.*

Proof. It is a trivial consequence of [5, Theor. 3.7] because the assumptions imply K -causality which is equivalent to stable causality. $\square$

The next result slightly generalize previous results due to Dowker, Garcia and Surya [2, Prop. 2] and Hawking and Sachs [4, Theor. 2.1]. The proof that (i) and distinction⁴ implies (iv), originally given in [4, Theor. 2.1D], would be particularly simple assuming the equivalence between stable causality and K -causality in light of theorem 6.2.

Theorem 6.4. *The following conditions for the spacetime (M, g) are equivalent*

- (i) (M, g) is future (resp. past) reflecting.
- (ii) For every $x \in M$, $\uparrow I^-(x) = I^+(x)$ (resp. $\downarrow I^+(x) = I^-(x)$).
- (iii) For every $x \in M$, $\bar{J}^+(x) = K^+(x)$ (resp. $\bar{J}^-(x) = K^-(x)$).

Defined

- (iv) For every $x \in M$, $\bar{J}^+(x) = J_S^+(x)$ (resp. $\bar{J}^-(x) = J_S^-(x)$).

then (iv) $\Rightarrow$ (i), moreover if (i) holds in the past or future case and the spacetime is distinguishing then (iv) holds (I give the proof of the last statement assuming the equivalence between K -causality and stable causality). Finally, if (i)-(iii) hold in the past or future case, then $\bar{J}^+ = K^+$, while if (iv) holds in the past or future case then, $\bar{J}^+ = K^+ = J_S^+$.

Proof. (i) $\Leftrightarrow$ (ii). The spacetime is future reflecting iff for every $x \in M$, $A^+(x) = \bar{I}^+(x)$ that is iff $\uparrow I^-(x) = \bar{I}^+(x)$, that is iff $\uparrow I^-(x) = I^+(x)$.

(i) $\Rightarrow$ (iii). $\bar{J}^+$ is not only closed but also transitive because if $(x, y) \in \bar{J}^+$ and $(y, z) \in \bar{J}^+$ then [6, Prop. 3.45], $y \in \bar{J}^+(x)$ and $z \in \bar{J}^+(y)$ from which it follows because of [5, theorem 3.3] (or [2, Claim 1]), $(x, z) \in \bar{J}^+$, hence $K^+ = \bar{J}^+$. In particular $\bar{J}^+(x) = \{y \in M : (x, y) \in \bar{J}^+\} = \{y \in M : (x, y) \in K^+\} = K^+(x)$, where in the first equality we used the future reflectivity.

(iii) $\Rightarrow$ (i). $x \in \bar{J}^-(z) \Rightarrow (x, z) \in \bar{J}^+ \Rightarrow (x, z) \in K^+ \Leftrightarrow z \in K^+(x) \Leftrightarrow z \in \bar{J}^+(x)$, thus (M, g) is future reflecting.

(iv) $\Rightarrow$ (i). $x \in \bar{J}^-(z) \Rightarrow (x, z) \in \bar{J}^+ \Rightarrow (x, z) \in J_S^+ \Leftrightarrow z \in J_S^+(x) \Leftrightarrow z \in \bar{J}^+(x)$, thus (M, g) is future reflecting.

Distinction and (i) $\Rightarrow$ (iv). I give the proof assuming that K -causality coincides with stable causality. From (i), say future case, it follows (iii) in the future case. Moreover, from theorem 6.3 the spacetime is stably causal (K -causal) thus, by theorem 6.2, $K^+ = J_S^+$, hence the thesis.

The last statement follows from the reflectivity since in this case $\bar{J}^+ = \{(x, z) : z \in \bar{J}^+(x)\}$. $\square$

Remark 6.5. It can be $\bar{J}^+ = K^+$ and yet the spacetime can be non-reflecting. An example is provided by 1+1 Minkowski spacetime in which a spacelike geodesic segment has been removed. Thus $\bar{J}^+ = K^+$ does not imply (iii) whereas the converse holds.

⁴ Examples of non-distinguishing spacetimes for which (i) holds but (iv) does not hold are for instance examples 5.1 and 5.2.

7 Conclusions

The relationship between stable causality and K -causality and their possible equivalence has been studied in detail. To this end new results for the K^+ future have been obtained (lemma 5.13, theorems 5.15, 5.16, 5.17). Unfortunately, a proof of the equivalence has not been given. A partial result in the direction of the equivalence has been the proof that in a K -causal spacetime, chosen an event, the light cones can be widened in a neighborhood of the event without spoiling K -causality (theorem 5.19). The process of enlarging the light cones can be continued and thus, if the equivalence does indeed hold, a final proof could be perhaps be obtained through an inductive process starting from this result. In any case, if the equivalence holds, in a K -causal spacetime the K^+ future coincides with the Seifert future J_S^+ (theorem 6.2). If the spacetime is not K -causal one expects to find some examples which show that in general $K^+ \neq J_S^+$ and indeed I gave the example of a causal spacetime (example 5.2). Finally, a new proof that Seifert's causal relation is a partial order iff the spacetime is stably causal has been given which uses some new lemmas which seem interesting in their own right (lemmas 3.9 and 3.11).

Acknowledgements

I warmly thank F. Dowker, R. Low, R.D. Sorkin and E. Woolgar, for sharing their impressions on the possible equivalence between stable causality and K -causality.

References

- [1] E. Aguirre-Dabán and M. Gutiérrez-López. On the topology of stable causality. *Gen. Relativ. Gravit.*, 21:45–59, 1989.
- [2] H. F. Dowker, R. S. Garcia, and S. Surya. K -causality and degenerate spacetimes. *Class. Quantum Grav.*, 17:4377–4396, 2000.
- [3] S. W. Hawking and G. F. R. Ellis. *The Large Scale Structure of Space-Time*. Cambridge University Press, Cambridge, 1973.
- [4] S. W. Hawking and R. K. Sachs. Causally continuous spacetimes. *Commun. Math. Phys.*, 35:287–296, 1974.
- [5] E. Minguzzi. The causal ladder and the strength of K -causality. I *Class. Quantum Grav.* 2007, In press.
- [6] E. Minguzzi and M. Sánchez. *The causal hierarchy of spacetimes*. Cont. to Proc. of the ESI Semester ‘Geometry of pseudo-Riemannian Manifolds with Application to Physics’, ed. D. Alekseevsky and H. Baum (ESI, Vienna, Sept-Dec 2005) (European Mathematical Society Publishing House), gr-qc/0609119, To appear.

- [7] B. O'Neill. *Semi-Riemannian Geometry*. Academic Press, San Diego, 1983.
- [8] R. Penrose. *Techniques of Differential Topology in Relativity*. Cbms-Nsf Regional Conference Series in Applied Mathematics. SIAM, Philadelphia, 1972.
- [9] I. Rácz. Causal boundary for stably causal spacetimes. *Gen. Relativ. Gravit.*, 20:893–904, 1988.
- [10] H. Seifert. The causal boundary of space-times. *Gen. Relativ. Gravit.*, 1:247–259, 1971.
- [11] R. D. Sorkin and E. Woolgar. A causal order for spacetimes with C^0 Lorentzian metrics: proof of compactness of the space of causal curves. *Class. Quantum Grav.*, 13:1971–1993, 1996.